



Mermaid syndrome, Sirenomelia: A case report and review of literature

Anahita Nosrati¹
Farshad Naghshvar²
Zhila Torabizadeh^{3*}
Omid Emadian⁴

^{1,2,3,4}Department of Pathology, Faculty of Medicine, Mazandaran University of Medical Science, Sari, Iran.

ARTICLE INFO

Article type:

Case and Review

Article history:

Received: 15 Sep 2012

Revised: 29 Dec 2012

Accepted: 8 Jan 2013

Keywords:

Sirenomelia, Mermaid syndrome, Oligohydramnios, Dextrocardia

<http://jpr.mazums.ac.ir>

ABSTRACT

Sirenomelia is a rare anomaly of caudal region of the body presented with fusion of lower limbs in which Genito-urinary, Gastrointestinal, Cardiovascular and Neural tube anomalies are found in most cases. We introduce a case of severe oligohydramnios, which had characteristic aspects of Sirenomelia at the time of delivery and review the pertinent articles. All in all it is our aim to present another case of sirenomelia to add to existing knowledge and data about the condition in order to make an earlier diagnosis and put an end to pregnancy safely in advance.

Introduction

Sirenomelia or Mermaid syndrome was originally stated by Rocheas in 1542 and Palfya in 1553 and named after the mythical Greek Siren.^{1,2,3} For the first time, Homer explained in the tempting singing creatures that seduced sailors to their deaths.⁴ Historical texts of around 2500 BC found in several European countries included references to female hybrids.⁴ It occurs about 1 in 100,000 live births.⁵ The most striking yet inconstant view of this malformative disorder is the complete or partial fusion of the lower limbs. It is a congenital anomaly including renal agenesis, ambiguous external genitalia, imperforate anus and blind intestinal loop.^{6,7} Occasionally

double inferior Vena cava and angiomatous lumbosacral myelocystocele are reported as well.^{8,9} Most of the Sirenomelia come to an end as stillbirth.⁶ Only a few numbers of whom have cited to be alive after delivery.^{10,11} Interestingly it has been said that there is an association with maternal diabetes.^{1,12} Hence we aim to present a fetus with features of mermaid syndrome in our department for first time and review proper articles.

Case presentation

A 34 years old primigravid pregnant woman at 25 weeks of gestation is counseled to terminate her pregnancy because of marked

*Corresponding Author: Zhila Torabizadeh MD, Associate Professor of Pathology

Mailing Address: Department of pathology, Emam Hospital, Amir Mazandarani boulevard, Sari, Iran

Tel: +98 151 2261701-4

Email: zhtorabi@yahoo.com

oligohydramnios that was incompatible with life. She had undergone an Ultrasound examination at 10 weeks gestation without apparently any reported abnormality in fetus. There were no medical or obstetric risk factors. There was no personal or family history for Diabetes Mellitus or congenital anomalies in relatives.

Routine Ultrasound examination, at 24 weeks of gestation revealed a live single fetus corresponding to 21 weeks gestation, weight: 270 gram. There were no single measurable pocket of amniotic fluid (severe Oligohydramnios); moreover, fetal kidneys, urinary bladder and stomach were not visualized.

A repeat Ultrasound was done in another institute and it turned out to be quite dextrocardia and left diaphragmatic hernia. Since there was severe oligohydramnios, the sonographer was not able to precisely assess both lower limbs and lumbar spines. Eventually after couple's consent termination is done at 25 weeks.

During autopsy procedure, fusion of the whole lower limbs from the hips to the ankles was founded out (Figure-1). In addition, it was realized syndactyly of the left foot with soft tissue union of the first and second toes.

The head was compressed and showed low set ears.

The face displayed bilateral complete lip and palate clefts. The lumbar spine showed a defect at the level of $L_3 - L_4$. There was no anal opening and no discernable genitalia.

The umbilical stump revealed just one artery and one vein. Complete absence of genitourinary system was evident. Kidneys, ureters and bladder were absent too. There was no stomach, and the midgut loop was blindly ended. The heart was located in the right side of the chest cavity (dextrocardia) (Figure-2) The lungs were non-flatted. The mentioned findings

were consistent with the diagnosis of sirenomelia.



Figure 1. Mermaid fetus, showing fused lower limbs and lip and palate clefts

Microscopic examination revealed just partially autolysed organs and generally not special pathologic changes.



Figure 2. Mermaid fetus, Dextrocardia

Discussion

Sirenomelia is a rare and lethal congenital anomaly with unknown etiology showing abnormal development of the caudal region of the body involving varying degrees of fusion of the lower limbs with or without bony defects (Table-1).

Other visceral defects such as hypo plastic lungs, cardiac agenesis, absent genitalia, digestive defects, absent kidney and bladder,

Table 1. Antenatal cases of mermaid syndrome diagnosed in the first trimester

Author	Year of publication	Number of references
Duhamel B	1961	26
Von Lennep E	1985	19
Steven RE, Hibelink DR	1986	17,18
Stocker JT	1987	27
Sirtori M	1989	21
Tang TT, Rodriguez JI	1991	13,14
Sarpong S, Kallen B	1992	20,5
Lynch SA	1995	24
Jaiyesimi F	1998	23
Aslan H	2001	12
Taori KB , Stanton MP	2002	7,11
Browne M	2004	9
Pinette MG	2005	15
Bakhtar O, Romano S, Vankeirsblick j, Ladure H	2006	22,4,2,3
Akbayir O	2008	29
Sikandar R	2009	1
Fadhlaoui A	2010	25
Ugwa RO	2011	6
Our current case	2012	-

vertebral and central nervous system defect are also reported.^{13, 14} The poor prognosis is usually forecasted depending on some kind of the visceral anomalies. Death usually stems from renal agenesis or dysgenesis¹⁵, which there likely seems to be happened to our case.

Risk factors

Many different kinds of theories have been suggested such as maternal diabetes^{6, 7, 8, 16}; however, our mother was not known to be diabetic. Genetic and environmental factors are also noted.^{7, 17} Hibelink et al pointed out that teratogenic agents like cadmium element and lead may cause sirenomelia in the golden hamster.¹⁸ The likely potential teratogenic effect of vitamin A is mentioned by Vonlennep et al¹⁹ and cocaine²⁰ or irradiation exposure²¹ as well. Association of the sirenomelia with new reproductive technologies, namely with ICSI (Intra Cytoplasmic Sperm Injection) was also described.²²

Besides, it has been reported in twins. Reports indicate a 100–150 times higher incidence in monozygotic twins relative to dizygotic twins or singletons.²³ In our patient, no twin gestation or family history of twinning was found out.

Another huge concern is genetic transmission. Lynch *et al*²⁴ realized an autosomal type of caudal dysgenesis, but no chromosomal defects are discovered in mermaid syndrome. This was an encouraging view for our mother to have sort of genetic consultation. Genetic counseling should even be suggested because the mentioned risk of recurrence is fairly 3–5%.²⁵ All abnormalities in our case like imperforate anus, a neural tube defect, limbs abnormality, cardiac dextrocardia and renal agenesis are frankly compatible with concept of sirenomelia in other similar related articles as well.²⁵

Pathogenesis

The pathogenesis of this syndrome has been discussed over time. Numerous theories have

been proposed to explain its origin. Five pathogenetic theories of sirenomelia are described:

1. An embryological insult¹⁵
2. Vascular steal theory¹⁷
3. As part of the caudal regression syndrome (CRS)²⁶
4. As part of the VACTERL syndrome (vertebral defects, anal atresia, cardiac defects, tracheo-esophageal fistula, renal anomalies, and limb abnormalities)²³
5. External forces acting on the caudal extremity.²⁵

From an embryological point of view, the sirenomelia sequence comes from an 'caudal mesoderm damage taking place between days 28–32 of fetal life (first theory).¹⁵ Therefore there are renal agenesis, lack of genital organs, imperforated anus, vertebral dysgenesis, and lower limb atrophy.²⁵

Second theory, Sirenomelia is probably a consequence of an abnormal blood supply of caudal parts of the embryo. A diverged abnormal vessel carries blood through the umbilical cord into the placenta. The vessel "steals" blood from the structures located below its origin from the tissues of fetal caudal part.¹⁶

A third theory regards sirenomelia as part of the caudal regression syndrome (CRS), a rare congenital anomaly described by Duhamel²⁶; including sacrum dysgenesis, spinal cord defect, urinary incontinence and misplaced lower limbs. Some inconstant features are renal dysgenesis and imperforate anus.

According to the fourth theory syndrome in the literature, sirenomelia could be a member of the VACTERL syndrome. VACTERL syndrome consists of vertebral, anal, cardiovascular, trachea-esophageal fistula, renal and limb dysgenesis. A major overlap has been realized in the phenotypic features of sirenomelia and VACTERL.²³

A fifth theory is the pressure theory stating hypoplasia of caudal extremity is owing to

external forces.²⁵ However not accepted on the whole.²⁵ Other less appropriate theories have also been proposed. Yet the overlap in these syndromes/theories is a matter of debate in the scientific world as to the uniqueness or varieties of these syndromes.

Classification

Stocker and Heifetz²⁷ classified the sirenomelia sequence into 7 types as below:

1. All thigh and leg bones are present
2. Single fibula
3. Absent fibula
4. partially fused femurs fused fibulae
5. partially fused femurs
6. single femur single tibia
7. single femur absent tibia

We did not have any radiographs of the lower limbs, but our case turned out to be pretty type six.

Diagnosis

It has been reported the possible first trimester diagnosis of mermaid fetus.³ Although the pregnant mother was supervised, it didn't take place in our case. The use of color Doppler imaging and MRI is claimed to have beneficial influence on in advance assessment.^{1, 2, 28}

Prognosis

Sirenomelia is fatal in most cases because of the characteristic pulmonary hypoplasia and renal agenesis. In children borned alive death will occur rather within a couple of days following birth²⁷. Presence of kidney is mandatory to survive. Murphy et al²⁴ reported a survived sirenomelia neonate. Laparotomy and colostomy were done, and eventually lower extremities were separated. It stands to reason that its managing is tough and rather costly with unforeseeable results in the case described by Stanton.¹¹ The patient had five interventions

between the age of 15 days and 4 years (she has been bedridden and dependent).

Conclusion

Sirenomelia is a kind of scarce and deadly congenital malformation. As far as we are concerned it is pretty much uncommon condition, so giving elaborated information may provide clinicians with a great chance to terminate the pregnancy in advance much more safely.

Acknowledgements

We thank Dr K. khodadadi and Miss M. Rashidi who provided us with advanced medical search and technical support to reach our aim.

Conflict of Interest

None declared.

Funding/Support

None declared.

References

1. Sikandar R, Munim S. Sirenomelia, the Mermaid syndrome: case report and a brief review of literature. JPMA The Journal of the Pakistan Medical Association. 2009; 59(10): 721.
2. Van Keirsbilck J, Cannie M, Robrechts C, de Ravel T, Dymarkowski S, Van den Bosch T, et al. First trimester diagnosis of sirenomelia. Prenatal diagnosis. 2006; 26(8): 684-8.
3. Ladure H, D Herve D, Loget P, Poulain P. Prenatal diagnosis of sirenomelia. Journal de Gynecologie Obstetrique et Biologie de la Reproduction. 2006; 35(2): 181.
4. Romano S, Esposito V, Fonda C, Russo A, Grassi R. Beyond the myth: The mermaid syndrome from Homer to Andersen: A tribute to Hans Christian Andersen's bicentennial of birth. European journal of radiology. 2006; 58(2): 252-9.
5. Källén B, Castilla EE, Lancaster P, Mutchinick O, Knudsen L, Martínez-Frías M, et al. The cyclops and the mermaid: an epidemiological study of two types of rare malformation. Journal of medical genetics. 1992; 29(1): 30-5.
6. Ugwu RO, Eneh AU, Wonodi W. Sirenomelia in a Nigerian triplet: a case report. Journal of medical case reports. 2011; 5(1): 426.
7. Taori K, Mitra K, Ghonga N, Gandhi R, Mammen T, Sahu J. Sirenomelia sequence (mermaid): Report of three cases. Indian Journal of Radiology and Imaging. 2002; 12(3): 399.
8. Assimakopoulos E, Athanasiadis A, Zafarakas M, Dragoumis K, Bontis J. Caudal regression syndrome and sirenomelia in only one twin in two diabetic pregnancies. Clinical and experimental obstetrics & gynecology. 2004; 31(2): 151.
9. Browne M, Fitchev P, Adley B, Crawford SE. Sirenomelia with an angiomatous lumbosacral myelocystocele in a full-term infant. Journal of perinatology. 2004; 24(5): 329-31.
10. Sammons MB. Shiloh pepin: The mermaid girl. Available from: <http://www.aolhealth.com/health/mermaid-syndrome>. [Last accessed on 2009].
11. Stanton MP, Penington EC, Hutson JM. A surviving infant with sirenomelia (Mermaid syndrome) associated with absent bladder. Journal of pediatric surgery. 2003; 38(8): 1266-8.
12. Aslan H, Yanik H, Celikaslan N, Yildirim G, Ceylan Y. Prenatal diagnosis of Caudal Regression Syndrome: a case report. BMC pregnancy and childbirth. 2001; 1(1): 8.
13. Tang TT, Oechler HW, Hinke DH, Segura AD, Franciosi RA. Limb body-wall complex in association with sirenomelia sequence. American journal of medical genetics. 1991; 41(1): 21-5.
14. Rodríguez JI, Palacios J, Razquin S. Sirenomelia and anencephaly. American journal of medical genetics. 1991; 39(1): 25-7.
15. Pinette MG, Hand M, Hunt RC, Blackstone J, Wax JR, Cartin A. Surviving sirenomelia. Journal of ultrasound in medicine. 2005; 24(11): 1555-9.
16. Tanha FD, Googol N, Kaveh M. Sirenomelia (mermaid syndrome) in an infant of a diabetic mother. Acta Medica Iranica. 2003; 41(1): 69-72.
17. Stevenson RE, Jones KL, Phelan MC, Jones MC, Barr Jr M, Clericuzio C, et al. Vascular steal: the pathogenetic mechanism producing sirenomelia and associated defects of the viscera and soft tissues. Pediatrics. 1986; 78(3): 451-7.
18. Hilbelink DR, Kaplan S. Sirenomelia: Analysis in the cadmium- and lead-treated golden hamster. Teratogenesis, carcinogenesis, and mutagenesis. 2005; 6(5): 431-40.
19. Von Lennep E, El Khazen N, De Pierreux G, Amy J, Rodesch F, Van Regemorter N. A case of partial sirenomelia and possible vitamin A teratogenesis. Prenatal diagnosis. 1985; 5(1): 35-40.
20. Sarpong S, Headings V. Sirenomelia accompanying exposure of the embryo to cocaine. South Med J. 1992; 85(5): 545-7.
21. Sirtori M, Ghidini A, Romero R, Hobbins J. Prenatal diagnosis of sirenomelia. Journal of ultrasound in medicine. 1989; 8(2): 83-8.

22. Bakhtar O, Benirschke K, Masliah E. Sirenomelia of an intracytoplasmic sperm injection conceptus: a case report and review of mechanism. *Pediatric and Developmental Pathology*. 2006; 9(3): 245-53.
23. Jaiyesimi F, Gomathinayagam T, Dixit A, Amer M. Sirenomelia Without Vitelline Artery Steal. *Annals of Saudi medicine*. 1998; 18: 542-4.
24. Lynch SA, Bond PM, Copp AJ, Kirwan WO, Nour S, Balling R, et al. A gene for autosomal dominant sacral agenesis maps to the holoprosencephaly region at 7q36. *Nature genetics*. 1995; 11(1): 93-5.
25. Fadhlou A, Khrouf M, Gaigi S, Zhioua F, Chaker A. The Sirenomelia sequence: a case history. *Clinical medicine insights Case reports*. 2010; 3: 41.
26. Duhamel B. From the mermaid to anal imperforation: the syndrome of caudal regression. *Archives of Disease in Childhood*. 1961; 36(186): 152-5.
27. Stocker JT, Heifetz SA. Sirenomelia. A morphological study of 33 cases and review of the literature. *Perspectives in pediatric pathology*. 1987; 10: 7.
28. Akbayir O, Gungorduk K, Sudolmus S, Gulkilik A, Ark C. First trimester diagnosis of sirenomelia: a case report and review of the literature. *Archives of gynecology and obstetrics*. 2008; 278(6): 589-92.