



## Factors Contributing to Postanesthetic Emergence Agitation in Pediatric Anaesthesia

Soheila Shahmohammadi<sup>1</sup>  
Afshin Gholipour Baradari<sup>2</sup>  
Mohammad Reza Habibi<sup>3\*</sup>  
Mohammad Jafar Saffar<sup>4</sup>

<sup>1</sup>CRNA, Research Fellow, Mazandaran University of Medical Sciences, Sari, Iran

<sup>2,3</sup>Department of Anesthesiology, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

<sup>4</sup>Department of Pediatric Infectious Disease, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

### ARTICLE INFO

*Article type:*  
Review Article

*Article history:*  
Received: 25 May 2013  
Revised: 17 June 2013  
Accepted: 11 July 2013

*Keywords:*  
Emergence, Agitation, General Anaesthesia, Pediatric

<http://jpr.mazums.ac.ir>

### ABSTRACT

Emergence Agitation that has been first described by Eckenhoff et al. in 1960's is a dissociated state of consciousness in which the child is inconsolable, irritable, uncooperative, typically thrashing, crying, moaning or incoherent. It is also a common problem in pediatric postanesthetic care unit with an incidence ranging from 10 to 80%. This literature review focused on presence of Emergence Agitation and contributing factors in children under general anaesthesia. It was conducted on Medline in PubMed area, Alta Vista Data bases, CINHALL and Google scholar in January 2013 for publications written in English with the following keywords: "Emergence Agitation, Etiology, Treatment, Pediatric Anaesthesia, Postanesthetic Care Unit, Children, inhaled anesthetics, intravenous anesthetics and Post Anesthetic Emergence Delirium". In this paper, we intend to review the factors contributing postanesthetic emergence agitation in children to improve our vision in this area.

### Introduction

Nowadays, about four million children undergo anaesthesia annually and Emergence Agitation (EA) has been identified as a significant problem in children at Post Anaesthesia Care Unit (PACU).<sup>1</sup> Early epidemiologic studies

demonstrated a 5.3% incidence of EA in all postoperative patients, with a more frequent incidence in children (12-13%).<sup>1,2</sup> The incidence of EA in Children who received volatile anesthetic agents (sevoflurane and

\*Corresponding author: Mohammadreza Habibi, Assistant Professor, Anesthesiologist, Fellowship of Cardiac Anesthesiology

Mailing Address: Department of Anesthesiology, Fatemeh Zahra Cardiac Center Hospital, Artesh Boulevard, Sari, Iran

Tel: +98 151 2223023-25

Fax: +98 151 2228484

Email: mohammadreza.habibi@gmail.com

desflurane) has been reported from 24 to 66%,<sup>3</sup> increasing to 80% in preschool children.<sup>4</sup> EA is a postanesthetic problem that interferes with child's recovery and presents a challenging situation for the post-anaesthesia care provider in terms of assessment and management.<sup>3</sup> Although, several factors have been identified as etiologic factors of EA, there is no entire description for emergence agitation. Many different causes have been suggested, such as rapid awakening in an unfamiliar settings, painful events like surgical wounds, agitation on induction, airway obstructions, environmental disturbances, duration of anaesthesia, hyperthermia, hypothermia, type and site of operation, premedication, inhaled and intravenous anesthetics and the anesthetic technique.<sup>1,5-7</sup> Although EA is usually self-limited and occurs within the first 30-minutes of recovery in PACU, it can last up to 2 days and leads to physical damage, disconnected of intravenous catheters, removing of dressing or drainage tube and monitoring devices. On the other hand, controlling the agitated child needs more nursing care and more post-anaesthesia care providers. In addition, the administration of sedative and analgesic drugs is associated with increased recovery time and PACU discharge delay.<sup>8-10</sup> Generally, treatment in all cases mentioned above is directed to the correction of causative agents. Although, numerous medications have been studied to prevent or reduce EA in children, no special preventive method has been shown to be highly superior. Understanding the risk factors for EA helps us to determine the best way to control this phenomenon in the PACU period. Because of the contradictory results of previous studies conducted to determine the related risk factors of EA and the former review article written about EA in children in 2011 reviewed this phenomenon as a whole, herein, we reviewed

the contributing factors for EA and suggested interventions.

## Materials and Methods

A literature review about the possible causes of postoperative agitation in preschool children was conducted on Medline in PubMed area, Alta Vista Data bases, CINHALL and Google scholar in January 2013 for publications written in English with the following keywords: "Emergence Agitation, Etiology, Treatment, Pediatric Anaesthesia, Postanesthetic Care Unit, Children, inhaled anesthetics, intravenous anesthetics and Post Anesthetic Emergence Delirium (PAED). All articles written in English, focused on presence of EA and contributing factors in children under general anaesthesia from January 2011 to January 2013 were included. Publications were excluded if the anaesthesia technique was not general.

## Results

A total 12 related articles met the criteria of our search, 10 of them were randomized control trials, one case report and one article was case control. Table 1 shows the list of the papers evaluated and the summary of the repossessed data.

## Discussion

This review reveals that Emergence Agitation (EA) still remains a significant postanesthetic problem that interferes with the child's recovery and challenges the PACU care provider in terms of assessment and treatment. Considering the potential risk factors is important to appropriately differentiate and treat agitation in the pediatric PACU. We also have found evidences that some of anesthetics may lead to decrease the incidence of postanesthetic EA.

**Table 1.** Characteristics and results of the studied papers (Continued)

Source /Date	Study Design	Cases	Age of patients	Anesthetic Technique	Premedication	Induction	Maintenance	Type of surgery	Agitation scale	EA <sup>c</sup> incidence
Na et al. 11 (2013)	RCT <sup>i</sup>	84	Pre-school children	Sevoflurane Anesthesia		thiopental, rocuronium, and 1% Sevoflurane	1% sevoflurane, 60% nitrous oxide in oxygen, and a continuous infusion of remifentanyl			remifentanyl group= 6 (4.25–10.25)
				Remifentanyl group (42 pt. <sup>a</sup> )				Adenotonsilectomy	PAED <sup>b</sup> & four-point EA scale	
Li et al. 12 (2013)	RCT	80	Pre-school children	Sevoflurane Anesthesia		thiopental, rocuronium, and 8% sevoflurane	2–3% sevoflurane			sevoflurane group= 11 (7.75–14.0)
				Sevoflurane Anesthesia (42 Pt.)				repair of unilateral inguinal hernia		

**Table 1.** Characteristics and results of the studied papers (Continued)

Source /Date	Study Design	Cases	Age of patients	Anesthetic Technique patient (No.)	Premedication	Induction	Maintenance	Type of surgery	Agitation scale	EA <sup>c</sup> incidence
<b>Kim et al.<sup>13</sup> (2013)</b>	RCT	222	18-72 months	Sevoflurane Anesthesia propofol 1 mg kg(-1) (Group P) at the end of Op.		Sevoflurane anaesthesia 8% + O2	Sevoflurane 2-2.5% + O2 50%	inguinal hernia repair	PAED	Group P < Group F < Group S
				Sevoflurane Anesthesia fentanyl 1 µg kg(-1) (Group F) at the end of Op.		Sevoflurane anaesthesia 8% + O2	Sevoflurane 2-2.5% + O2 50%			
<b>Chen et al.<sup>14</sup> (2013)</b>	RCT	84	2-7 y/o	Sevoflurane Anesthesia dexmedetomidine 1 µg.kg(-1) iv plus a 1 µg.kg(-1).hr(-1) infusion,		sevoflurane anaesthesia	sevoflurane	strabismus surgery	PAED	PAED scores for EA were lower in the dexmedetomidine (P < 0.001) and ketamine (P = 0.002) groups than in the placebo group
				Sevoflurane Anesthesia ketamine 1 mg.kg(-1) iv plus a 1 mg.kg(-1).hr(-1) infusion,		sevoflurane anaesthesia	sevoflurane			
				Sevoflurane Anesthesia Normal saline						



**Table 1.** Characteristics and results of the studied papers (Continued)

Source /Date	Study Design	Cases	Age of patients	Anesthetic Technique	Premedication	Induction	Maintenance	Type of surgery	Agitation scale	EA <sup>c</sup> incidence
<b>Dahmani et al.<sup>17</sup> (2012)</b>	Case report	1	3y/o		-	sevoflurane 6% (in a mixture of O <sub>2</sub> /N <sub>2</sub> O: 50%/ 50%)	spontaneously breathing with a 3% sevoflurane end-tidal concentration (in a mixture of O <sub>2</sub> /N <sub>2</sub> O:50% /50%)	bilateral myringotomy	PAED	PAED score=19
						Desflurane Anesthesia: group C received normal saline				
<b>Jeong et al.<sup>18</sup> (2012)</b>	RCT	60	2-8y/o		Atropine 0.01 mg/kg was injected intramuscularly 30 min before the induction of anesthesia	Thiopental sodium 5 mg/kg and rocuronium 0.6 mg/kg	oxygen 1.5 L/min, nitrous oxide 1.5 L/min and desflurane at 4-6 vol%	brief ophthalmic surgery	EA and the modified Children's Hospital of Eastern Ontario Pain Scale	K <sup>1</sup> .0<K0.5<C
						Desflurane Anesthesia: group K0.5 received ketamine 0.5 mg/kg 10 min before the end of the surgery				



**Table 1.** Characteristics and results of the studied papers (Continued)

Source /Date	Study Design	Cases	Age of patients	Anesthetic Technique patient (No.)	Premedication	Induction	Maintenance	Type of surgery	Agitation scale	EA <sup>c</sup> incidence
<b>LI et al.</b> <sup>21</sup> (2011)	RCT	105	3–11 y/o	Sevoflurane Anaesthesia: normal saline (control group), ----- Sufentanil 0.2 µg/kg (S2) ----- Fentanyl 2 µg/kg (F2) 1 minute after loss of the eyelash reflex	-	sevoflurane	sevoflurane	adenotonsilectomy	PAED scales	The incidence of severe agitation was significantly lower in S2 and F2 groups vs. the control group
<b>Lee et al.</b> <sup>22</sup> (2011)	RCT	56		Sevoflurane Anaesthesia: ET-A group (n = 56, endotracheal tube and extubation whilst awake) ----- Sevoflurane Anaesthesia: ET-D group (n=56, endotracheal tube and deep extu- bation) ----- Sevoflurane Anaesthesia: LMA-D group (n = 56, experienced LMA and deep removal)		8% sevoflurane in nitrous oxide/oxyge n (3/1 l/min) mixture via a face mask 2 µg/kg fentanyl for analgesia and 0.1 mg/kg ondansetron to prevent nausea and vomiting	sevoflurane with air/oxygen (1/1 l/min) mixture in a semiclosed circle system	Subumbilical Surgery	-	LMA-D <sup>g</sup> patients compared with patients in the ET-A <sup>h</sup> Group (21.4% Vs. 41.1%)

**Abbreviations:** <sup>a</sup> Pt (Patient), <sup>b</sup> PAED (Post anesthetic Emergence Delirium), <sup>c</sup> EA (Emergence Agitation), <sup>d</sup> NS (No Significant), <sup>e</sup> VAS (visual analog scale), <sup>f</sup> K (Ketamine), <sup>g</sup> Laryngeal Mask Airway – Deep (LMA-D), <sup>h</sup> Endotracheal- Awake, <sup>i</sup> Randomized Clinical Trial.



Na et al. (2013) in a study to investigate the effect of sevoflurane anaesthesia in combination with remifentanyl during the induction and maintenance of anaesthesia found that PAED score in remifentanyl group was significantly lower than sevoflurane group ( $P=0.007$ ) and the proportion of patients with PAED scores 10 and four point scale scores 3 were significantly lower in the remifentanyl group than in the sevoflurane group.<sup>11</sup> Comparing the effect of sufentanil and fentanyl by Li et al. on emergence agitation in preschool children who underwent repair of unilateral inguinal hernia following sevoflurane anaesthesia have showed that  $0.15\mu\text{g}/\text{kg}$  sufentanil compared with a single dose of  $1.5\mu\text{g}/\text{kg}$  fentanyl could significantly decrease the incidence of emergence agitation without delaying the recovery time.<sup>12</sup>

Similarly, in a study by Lee et al., to investigate the effect of sufentanil to reduce emergence agitation after sevoflurane anaesthesia in children undergoing adenotonsillectomy compared with fentanyl, they had concluded that administration of sufentanil at  $0.2\mu\text{g}/\text{kg}$  after induction of anaesthesia could reduce emergence agitation more without delaying the recovery time or causing significant hypotension in children compared with fentanyl.<sup>21</sup>

It has been shown that propofol comparing with fentanyl to prevent of EA after sevoflurane anaesthesia in children was more effective and associated with lower PAED score.<sup>13</sup>

Intraoperative administration of ketamine and dexmedetomidine has also decreased the incidence of EA and PAED score in pediatric patients under sevoflurane anaesthesia.<sup>14</sup>

The incidence of EA has been reported higher with sevoflurane compared with desflurane and isoflurane. However, among the three anesthetic agents no correlation was found between the incidence of EA and duration of anaesthesia or age.<sup>15</sup> Evaluating the efficacy and safety of

dexmedetomidine for emergence agitation after tonsillectomy under sevoflurane anaesthesia in children appeared that dexmedetomidine could be safe and effective to reduce the incidence of early emergence agitation in children after tonsillectomy. Initial loading dose of  $1.0\mu\text{g}/\text{kg}$  followed by a maintenance infusion of  $0.4\mu\text{g}/\text{kg}/\text{hrs}$  was a better choice for children who received sevoflurane anaesthesia.<sup>16</sup>

Allowing one of the parents to enter the PACU and holding the child was associated with reduced risk factor of EA and/ or to treat it.<sup>17</sup>

Similarly, the results of Zand et al.'s study to compare the effects of midazolam premedication and parental presence during sevoflurane and halothane anaesthesia induction on the incidence of postoperative agitation in pediatric patients revealed that the presence of a parent at induction of sevoflurane anaesthesia was as effective as midazolam premedication in decreasing the incidence of postoperative agitation. Midazolam premedication effective to decrease postoperative EA associated with halothane was used as the anesthetic agent.<sup>20</sup>

The results of the study with the different dosages of ketamine with desflurane anaesthesia by Jeong et al. for brief ophthalmic surgery demonstrated that both the incidence of EA and pain scales were at least in K1.0 group compared with the K0.5 and placebo groups.<sup>18</sup>

Investigating the effect of remifentanyl as an alternative to N<sub>2</sub>O by Choi et al. in 2011 on EA and the presence of postoperative pain in preschool children under general anaesthesia with sevoflurane for tonsillectomy/adenoidectomy surgery indicated that severity post-operative pain in remifentanyl group was more than the N<sub>2</sub>O group ( $P=0.012$ ). There were no significant differences between the two groups in incidence of EA.<sup>19</sup>

Lee et al. in their study to compare the effect of laryngeal mask airway (LMA) and the removal of the LMA in a deeply anaesthetized state with endotracheal tube (ET) and extubation when the

patient was awake or deeply anaesthetized on the incidence of emergence agitation in preschool children after sevoflurane anaesthesia for subumbilical surgery concluded that using an LMA and deep removal could decrease postoperative emergence agitation more than endotracheal tube and awake extubation after sevoflurane anaesthesia in pediatric patients.<sup>22</sup>

## Conclusions

This review has identified that short time to awakening, sevoflurane anaesthesia, otorhinolaryngology procedures, preschool children age and difficult parental-separation behavior were the risk factors of EA. On the other hand the combination of remifentanyl, sufentanyl, propofol, ketamine and dexmedetomidine with sevoflurane anaesthesia and using an LMA and deep removal of ET reduce the incidence of EA.

## Conflict of Interest

None declared.

## Funding/Support

None declared.

## References

1. Nasar V.G., Hannallah R S. Emergence Agitation in children: A Review. *M.E.J. Anesth* 2011; 21(2): 175-184.
2. Eckenhoff JE, Kneale DH, Dripps RD: The incidence and etiology of postanesthetic excitement. *Anesthesiology* 1961; 22: 667-73.
3. Voepel-Lewis T, Malviya S, Tait AR. A prospective cohort study of emergence agitation in the pediatric postanaesthesia care unit. *AnesthAnalg* 2003; 96(6):1625-30.
4. Bortone L, Ingelmo P, Grossi S, Grattagliano C, Bricchi C, Barantani D, et al. Emergence agitation in preschool children: double-blind, randomized, controlled trial comparing sevoflurane and isoflurane anaesthesia. *PaediatrAnaesth* 2006; 16: 1138-43.
5. Beskow A, Westrin P. Sevoflurane causes more postoperative agitation in children than does halothane. *ActaAnaesthesiolScand* 1999; 43:536-41.
6. Hollister GR, Burn JM. Side effects of ketamine in pediatric anaesthesia. *AnesthAnalg* 1974; 53:264-7.
7. Brown R. Postoperative Recovery. In: Nagelhout J.J., Zagalaniczny K.L. *Nurse Anaesthesia*. 3<sup>rd</sup> Ed. USA: Elsevier Sunders 2005: 1148-1149.
8. Nicholau D. Postanaesthesia Recovery. In: Stoelting R.K., Miller R.D. *Basic of Anaesthesia*, fifth Ed. USA: Churchill Livingstone Elsevier 2007: 572-573.
9. Olympio MA. Postanesthetic delirium: historical perspectives. *J ClinAnesth* 1991; 3: 60-3.
10. Veyckemans F. Excitation phenomena during sevoflurane anaesthesia in children. *CurrOpinAnaesthesiol* 2001; 14:339-43.
11. Na HS, Song IA, Hwang JW, Do SH, Oh AY. Emergence agitation in children undergoing adenotonsillectomy: a comparison of sevoflurane vs. sevoflurane-remifentanyl administration. *ActaAnaesthesiolScand* 2013; 57(1):100-5.
12. Li X, Zhang Y, Zhou M, Xia Q, Li W, Lu Q. The effect of small dose sufentanyl on emergence agitation in preschool children following sevoflurane anaesthesia for elective repair of unilateral inguinal hernia. *Saudi Med J* 2013; 34(1):40-5.
13. Kim MS, Moon BE, Kim H, Lee JR. Comparison of propofol and fentanyl administered at the end of anaesthesia for prevention of emergence agitation after sevoflurane anaesthesia in children. *Br J Anaesth* 2013; 110(2):274-80.
14. Chen JY, Jia JE, Liu TJ, Qin MJ, Li WX. Comparison of the effects of dexmedetomidine, ketamine, and placebo on emergence agitation after strabismus surgery in children. *Can J Anaesth* 2013; 60(4): 385-92.
15. Singh R, Kharbanda M, Sood N, Mahajan V, Chatterji C. Comparative evaluation of incidence of emergence agitation and post-operative recovery profile in paediatric patients after isoflurane, sevoflurane and desflurane anaesthesia. *Indian J Anaesth* 2012; 56(2):156-61.
16. Meng QT, Xia ZY, Luo T, Wu Y, Tang LH, Zhao B, Chen JH, Chen X. Dexmedetomidine reduces emergence agitation after tonsillectomy in children by sevoflurane anaesthesia: a case-control study. *Int J Pediatr Otorhinolaryngol* 2012; 76(7): 1036-41.
17. Dahmani S, Mantz J, Veyckemans F. Case scenario: severe emergence agitation after myringotomy in a 3-yr-old child. *Anesthesiology* 2012; 117 (2): 399-406.
18. Jeong WJ, Kim WY, Moon MG, Min DJ, Lee YS, Kim JH, Park YC. The effect of ketamine on the separation anxiety and emergence agitation in children undergoing brief ophthalmic surgery under

- desflurane general anaesthesia. *Korean J Anesthesiol* 2012; 63 (3): 203-8.
19. Choi HR, Cho JK, Lee S, Yoo BH, Yon JH, Kim KM. The effect of remifentanil versus N<sub>2</sub>O on postoperative pain and emergence agitation after pediatric tonsillectomy/adenoidectomy. *Korean J Anesthesiol* 2011; 61(2):148-53.
20. Zand F, Allahyary E, Hamidi AR. Postoperative agitation in preschool children following emergence from sevoflurane or halothane anaesthesia: a randomized study on the forestalling effect of midazolam premedication versus parental presence at induction of anaesthesia. *Acta Anaesthesiol Taiwan* 2011; 49(3):96-9.
21. Li J, Huang ZL, Zhang XT, Luo K, Zhang ZQ, Mao Y, et al. Sufentanil reduces emergence agitation in children receiving sevoflurane anaesthesia for adenotonsillectomy compared with fentanyl. *Chin Med J (Engl)* 2011; 124(22): 3682-5.
22. Lee YC, Kim JM, Ko HB, Lee SR. Use of laryngeal mask airway and its removal in a deeply anaesthetized state reduces emergence agitation after sevoflurane anaesthesia in children. *J Int Med Res.* 2011; 39(6):2385-92.