



## Vitamin E in children with asthma: A review

Javad Ghaffari<sup>1\*</sup>

Hossein Ashrafi<sup>2</sup>

Ali Reza Ranjbar<sup>3</sup>

Zeinab Nazari<sup>4</sup>

<sup>1</sup>Antimicrobial Resistant Nosocomial Infection Research Center, Mazandaran University of Medical Sciences, Sari, Iran

<sup>2</sup>Beatson Institute for Cancer Research, Faculty of Medicine, University of Glasgow, UK

<sup>3</sup>Research Institute of Interventional Allergology and Immunology, Bonn/Cologne, Germany

<sup>4</sup>Department of Obstetric & Gynecologist, Mazandaran University of Medical Sciences, Sari, Iran

### ARTICLE INFO

#### Article type:

Review Article

#### Article history:

Received: 18 June 2014

Revised: 22 Jul 2014

Accepted: 2 Aug 2014

#### Keywords:

Vitamin E, Asthma, Child

<http://jpr.mazums.ac.ir>

### ABSTRACT

Asthma is a chronic inflammatory airway disorder and the most common chronic disease among children characterized by frequent and chronic cough, wheezing, and dyspnea. Asthma is defined as an inflammatory process in response to T-Helper 2 cells. Vitamin E as a Lipid soluble vitamin; containing 4 isomers  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ , with  $\alpha$ -tocopherol as the most important isomer is an important defensive agent against epithelial tissue damages caused by antioxidants in human. The results of animal studies showed vitamins C and E cause decreased and modulation of pulmonary responses to oxidant agents such as O<sub>3</sub> or NO<sub>2</sub>. A few studies showed an association between reduced intake of vitamin E by mothers during pregnancy and wheezing in children aged 2 years old. Some studies showed significant declines in serum vitamin E level in asthmatic children compared to non-asthmatic. However, in another study no significant relationship was observed between serum level of vitamin E and risk of asthma. There are controversies on the effect of vitamin E in children with asthma. This study aimed to investigate the role of vitamin E in children with asthma.

### Introduction

Asthma is a chronic inflammatory airway disorder and the most common chronic disease among children characterized by

frequent and chronic cough, wheezing and dyspnea.<sup>1</sup> The prevalence of asthma is different worldwide with an increasing trend

\*Corresponding Author: Javad Ghaffari MD, Associate professor of allergy and clinical immunology, Mailing Address: Department of pediatric Immunology and Allergy & Antimicrobial Resistant Nosocomial Infection Research Center, Bou Ali Sina Hospital, Pasdaran Boulevard, Sari, Iran  
Tel: +98 11 33332331  
Fax: +98 11 33334506  
Email: javadneg@yahoo.com

in recent decades which imposes considerable financial and sociocultural burden on patients' families. More than 300 million patients are suffering from asthma around the world. The exact cause of asthma is not known yet. But, multi-factorial causes, such as genetic and environmental factors are involved.<sup>2,3</sup> Also, asthma is a complex disease that multiple factors such as autonomic system, immunology, infections, endocrine and psychological agents play an important role in the development of the disease.<sup>4</sup> Common environmental factors are infections, different allergens and pollutants that cause inflammation of respiratory tract and respiratory symptoms by effect on cell mediated immunity and producing different cytokines. Asthma is defined as an inflammatory process in response to T-Helper (TH2) cells. Cytokines like IL4, 5 and 13 that produce IgE can lead to realizing of mediators.<sup>5</sup>

For instance, viral agents increase production of fibroblast, neutrophils, eosinophils and monocytes by increasing Inter Cellular Adhesion Molecule 1 (ICAM-1) via nuclear factor kappa-light-chain-enhancer of activated B cells (NK-F $\beta$ ) and stimulating TH2 cells and increasing production of cytokine 4,6,8 and 16.<sup>3</sup> In recent years, there has been increasing interests in the role of nutrition in severity of asthma. This interest is more focused on the role of antioxidants such as vitamins A, C, E, and carotenoids. Some reports on development of asthma signified the role of insufficient levels of antioxidants such as vitamins and minerals following low intake of green plants (thus reduced level of vitamin E) and increase in consumption of foods containing free radicals.<sup>6-9</sup> Peripheral blood inflammatory cells and Bronchoalveolar lavage (BAL) fluid produce more anion superoxide in asthmatic

patients compared to control group which led to increasing of antioxidant consumption.<sup>10-12</sup> Steroids reduce oxygen free- radicals produced by these cells.<sup>13</sup> These antioxidants agents cause respiratory tract smooth muscle contraction, increasing of secretion, and sheering of epithelium. Reactive oxidant species (ROS) result in production of cytokines and chemokines.<sup>14, 15</sup> which proves the role of nutrition in development and severity of asthma. Vitamin E was used to decrease the symptoms of asthma and improving respiratory status of asthmatic patients.<sup>6</sup> It decreased or prevented toxicity of oxidant agents on respiratory system.<sup>6, 16-18</sup> There is a paucity of information regarding the effect of vitamin E on pediatric asthma, therefore, this study aimed to review the effect of vitamin E on pediatric asthma.

### **Vitamin E**

Vitamin E is the major Lipid soluble vitamin which is also an important defensive agent against epithelial tissue damages caused by antioxidants in human. It contains 4 isomers including  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ , amongst which  $\alpha$ -tocopherol is the most important isomer. This vitamin causes break down of pre-oxidase chain of fat and also decreasing production of prostaglandin E2 following peroxidation of fat in pulmonary epithelial cells after contact with Ozone (O<sub>3</sub>).<sup>19</sup> Antioxidant agents are the first defensive line against oxygen free-radicals. Fluid existence in respiratory tracts is containing antioxidants agents such as vitamin C, uric acid, glutathione peroxidase,  $\alpha$ -tocopherol (vitamin E), catalase, dismutase superoxide, thioredoxin reductase, seruloplasmin and transferrin. These agents are found in nutrients like fruits and vegetables.<sup>14, 16</sup> These agents have a role in clearance of free radicals and act as sacrificial targets for O<sub>3</sub>. They also restrict the interaction between oxidants with lipid and

proteins secreted by respiratory tract lining fluid.<sup>20</sup> Vitamin E is an important antioxidant in cell membrane containing tocopherol and tocotrienol that inhibits oxidation of components such as unsaturated fatty acid, even it can stimulate polyunsaturated fatty acid against antioxidants.<sup>4, 21</sup> In addition, this vitamin causes improvement of T-cells and granulocytes functions.<sup>21</sup> Oxidant are agents causing suppression of TH1 and increase in production of TH2. Previous studies showed an increase in the level of vitamin E in lung following consumption of vitamin E supplements compared to non-users. So that, vitamin E moves to the lung in oxidative stress situations.<sup>22</sup> vitamin E suppresses IgE response due to allergic agents during allergic diseases like asthma.<sup>23</sup> It also improves the blood supply and tissue repair.<sup>4</sup> Vitamin C was reported to be effective in clearance of free radicals in combination with vitamin E.<sup>4</sup> It seems vitamin E consumption is decreased in recent years. For example, in 1950 the intake of vitamin E was 13-15mg/day that decreased to 10.66mg/day in 2005- 2006. The cause of vitamin E deficiency is low intake of green vegetables, cereals and whole grains.<sup>24-</sup>  
<sup>26</sup> In children, vegetables may decrease wheezing.<sup>27</sup> However, a study in England revealed no considerable change in the prevalence and severity of asthma in school-age children who received fruits more than 1.3-2 serving/d.<sup>28</sup>

#### **Animal studies**

The results of animal studies showed the role of vitamins C and E in decreasing and modulation of pulmonary responses to antioxidant agents such as O<sub>3</sub> or NO<sub>2</sub>.<sup>29,30</sup> Vitamin E deficiency and decrease in levels of vitamin E lead to production of pro-inflammatory markers, decrease in surfactant lipid synthesis in type II alveolar cells and epithelial tissue damage due to air pollution.<sup>31</sup>

Elsayed showed that level of vitamin E in animal receiving vitamin E supplements was higher in pulmonary tissue after contact with Ozone than the animal which did not receive the supplements.<sup>32</sup>

#### **Human studies**

In another study mononuclear cells of cord blood in mothers of neonates who had higher intake of vitamin E during pregnancy demonstrated lower response to antigen stimulation compared to mothers with lower consumption of vitamin E.<sup>33</sup>

A study on children with asthma who were exposed to air pollution and received vitamin C and E supplements, found a considerable decrease in IL6 and IL8 level of nasal secretion compared to children who received placebo.<sup>34</sup> This result revealed the protective effect of antioxidant against the side effects of O<sub>3</sub> on pulmonary function in children with asthma suffering from mild vitamins deficiency in dietary intake.<sup>16</sup> In some controlled studies in human, using vitamin C and E supplements with different dosages caused protection against acute Ozone effects on lung function.<sup>35,36</sup>

In a study conducted by Romieu in children with asthma who received vitamin C 250mg/d combined with vitamin E 50mg/d, exposure to air pollution such as NO<sub>2</sub> and Ozone (particle mass less than 10 µm or PM<sub>10</sub>) showed the protective role of the supplements against acute Ozone effects on lung function. These effects were found to more in individuals with moderate asthma than those with mild asthma.<sup>16</sup> Other studies showed a relationship between intake of vitamin E and risk of asthma, and two even confirmed its protective effect.<sup>37-40</sup> A cohort study by devereux which was performed in 5-year-old children showed a reverse correlation between receiving vitamin E during pregnancy by their mothers and

wheezing and asthma as a present illness and the symptoms presented in one year ago.<sup>21</sup>

In a study by Nada Omar, there was a significant decline in serum vitamin E level in asthma children compared to non-asthmatic.<sup>4</sup> Similar findings were reported by others.<sup>41-43</sup> While, the result of another study was not similar.<sup>44</sup> The cause of different results reported by different studies could be due to several factors such as severity of asthma, diet and statistical techniques. For instance, Nada Omar indicated that decrease in levels of vitamin E in children was associated with more severe asthma which was reported by the others. Also, the levels of malondialdehyde (MDA) and ROS increased in asthmatic children.<sup>45, 46</sup>

The result of another study showed no significant relationship between serum level of vitamin E and risk of asthma. The serum level of vitamin E in children with asthma was not significantly different from that of the healthy individuals.<sup>47</sup> In a report, there was a significant relationship between decrease in consumption of vitamin E and FEF25-75. This finding is indicated that vitamin E plays a protective role on small respiratory tract function, but changing in FEV1 was not significant and was less sensitive.<sup>48</sup> An investigation in Saudi Arabia, showed that inadequate intake of vitamin E was associated with increased asthma attack.<sup>49</sup> Another study showed that severe asthma was associated with vitamin E intake, but it was not related to asthmatic status.<sup>50</sup> Several studies have been found correlations between higher intake of fruity vegetables, citrus and kiwi fruits and wheezing in children. Mediterranean foods due to rich in vegetables and fruits were associated with decreasing asthma and wheezing.<sup>51-54</sup> In another investigation Gilliland et al. showed that decreasing intake of vitamins A, C and E

was associated with decline of spirometry parameters including FVC, FEV1, FEF25-75.<sup>48</sup> Decreasing pulmonary function caused by low intake of vitamins was reported in other studies too.<sup>44,47</sup> Receiving high dose of antioxidants develop wheezing in children aged 2 years.<sup>55</sup> In a study by shock et al,  $\alpha$ -tocopherol level in BAL fluid was assessed in children with atopic asthma and healthy control group and no difference was found between the two groups. But, the results of other studies showed that the level of BAL fluid decreased in presence of gaseous pollutants such as Ozone, NO<sub>2</sub> and small particles,<sup>44,56-58</sup> Consumption of vegetables and fruits could decrease systemic inflammation specially decline in CRP concentration.<sup>59</sup> In pearson's study, vitamin E had no effect on pediatric asthma.<sup>18</sup> Other studies found that consuming antioxidant vitamins or vegetables and fruits improve pulmonary function or may decrease wheezing.<sup>60-63</sup>

#### **Study on mothers**

In a survey performed in USA, low consumption of vitamin by mothers during pregnancy was associated with wheezing in children aged 2 years old. Same results were also found in Japan, where increased level of wheezing was observed in children aged 16-24 months.<sup>64,65</sup>

Also, decreased  $\alpha$ -tocopherol (vitamin E) during pregnancy was associated with failure to thrive in pregnancy, decreased pulmonary function and increasing asthma in 5 year-old children.<sup>21</sup> But, a study showed high dose of vitamin E consumption in pregnant women with preeclampsia was not associated with decreasing wheezing and asthma in children.<sup>65</sup> In a study by Devereux, receiving vitamin E during pregnancy was associated with reduced proliferative response of mononuclear cells to allergens like grass and

mites compared to individuals who had low vitamin intake. Wheezing was also decreased in first group.<sup>33,66</sup> Similar results were also obtained for mites and cockroach.<sup>67</sup> Augusto in an investigation showed that higher antioxidants intake such as vitamin E and Zinc during pregnancy alleviates wheezing in children at 2 years of age, while it had no effect on eczema.<sup>55</sup>

Martindale et al. showed that consumption of vitamin E by mothers resulted in decreased level of wheezing in children aged 2 years old and atopic eczema.<sup>66</sup> In a study by Clark, the risk of asthma reduced in children whose mother received vitamin E 15mg/d during pregnancy.<sup>68</sup> In Anne's study, pregnant women at risk of preeclampsia, consumption of vitamin C and E had no effect on prevalence of asthma in their children aged 2 years old, unless, causes increasing cost and health care utilisation.<sup>65</sup>

## Conclusion

Although there were not many studies on the consumption of antioxidants such as vitamin E, vegetables and fruits in children with asthma and/ or pregnant women, but most of them indicated the positive effect of antioxidants intake such as vitamin E which found in most vegetables and fruits to relieve the prevalence of asthma or improving the symptoms of the disease and spirometry parameters among the patients.<sup>6,16,35-40</sup> However, some studies did not confirm this positive effect.<sup>47,48,50</sup> Therefore, further investigations and clinical trials are needed to confirm the effect of antioxidants such as vitamin E on asthma in children.

## Conflict of Interest

None declared.

## Funding/Support

None declared.

## References

1. Ghaffari J, Aarabi M. The prevalence of pediatric asthma in the Islamic Republic of Iran: A review and meta-analysis. *JPR* 2013; 1 (1): 2-11.
2. National Asthma Education and Prevention program. Expert Panel Report 2: Guidelines for the Diagnosis and Management of Asthma. Bethesda (MD): National Institutes of Health; 1997. National Institutes of Health publication No. 97-4051.
3. Farid Hossaini R, Ghaffari J, Ranjbar A, Haghshenas M, Rafatpanah H. Infections in Children with Asthma. *JPR* 2013; 1 (1) : 25-36.
4. Al-Abdulla NO, Al Naama LM, Hassan MK. Antioxidant status in acute asthmatic attack in children. *JPM* 2010; 60(12):1023-27.
5. Allan K, Devereux G. Diet and Asthma: Nutrition Implications from Prevention to Treatment. *J Am Diet Assoc* 2011; 111: 258-268.
6. Ghaffari J, Farid Hossaini R, Khalilian A, Nahanmoghdam N, Salehifar E, Rafatpanah H. Vitamin E Supplementation, Lung Functions and Clinical Manifestations in Children with Moderate Asthma: A Randomized Double Blind Placebo-Controlled Trial. *Iran J Allergy Asthma Immunol* 2014; 13(2):98-103.
7. National Food Survey datasets. UK Department of Environmental Farming and Rural Affairs. Website. <http://www.defra.gov.uk/evidence/statistics/foodfarm/food/familyfood/nationalfoodsurvey/index.htm>. Updated March 2010. Accessed December 9, 2010.
8. Ghaffari J, Rafatpanah H, Nazari Z, Abaskhanian A. Serum Level of Trace Elements (Zinc, Lead, and Copper), Albumin and Immunoglobulins in Asthmatic Children. *Zahedan Journal of Research in Medical Sciences*. 2013; 15(9): 27-30.
9. Murr C, Schroecksnadel K, Winkler C, Ledochowski M, Fuchs D. Antioxidants may

- increase the probability of developing allergic diseases and asthma. *Med Hypotheses*. 2005;64(5): 973-977.
10. Vachier I, Damon M, Le Doucen C, Crastes de Paulet A, Chanez P, Michel FB, Et al. Increased oxygen species generation in blood monocytes of asthmatic patients. *Am Rev Respir Dis* 146(5Pt1):1161-1166.
  11. Chanez P, Dent D, Yukawa T, Barnes PJ, Chung KF. Generation of oxygen free radicals from blood eosinophils from patients after stimulation with PAF or phorbol ester. *Eur Respir J*. 1990; 3(9):1002-7.
  12. Cluzel M, Damon M, Chanez P, Bousquet J, Crastes de Paulet A, et al. Enhanced alveolar cell luminol-dependent chemiluminescence in asthma. *J Allergy Clin Immunol* 80(2):195-201.
  13. Magori M, Vachier I, Godard P, Force M, Bousquet J, Chanez P. Superoxide anion production by monocytes of corticosteroid-treated asthmatic patients. *Eur Respir J* 1998; 11(1):133-138.
  14. Wood LG, Garg ML, Smart JM, Scott HA, Barker D, Gibson PG. Manipulating antioxidant intake in asthma: a randomized controlled trial. *Am J Clin Nutr*. 2012 Sep;96(3):534-43. doi: 10.3945/ajcn.111.032623. Epub 2012 Aug 1.
  15. Wood LG, Gibson PG, Garg ML. Biomarkers of lipid peroxidation, airway inflammation and asthma. *Eur Respir J* 2003; 21(1): 177-86.
  16. Romieu II, Sienna-Monge JJ, Ramírez-Aguilar M, Téllez-Rojo MM, Moreno-Macías H, Reyes-Ruiz NI, et al. Antioxidant supplementation and lung functions among children with asthma exposed to high levels of air pollutants. *Am J Respir Crit Care Med* 2002; 166(5):703-9.
  17. Gao J, Gao X, Li W, Zhu Y, Thompson PJ. Observational studies on the effect of dietary antioxidants on asthma: A meta-analysis. *Respirology* 2008; 13(4):528-36.
  18. Pearson PJK, Lewis SA, Britton J, Fogarty A. Vitamin E supplements in asthma: A parallel group randomized placebo controlled trial. *Thorax* 2004; 59(8): 652-6.
  19. Burton GW, Ingold KU. Autooxidation of biological molecules. 1. The antioxidant activity of vitamin E and related chain-breaking phenolic antioxidants in vitro. *J Am Chem Soc* 1981; 103(21): 6472-6477.
  20. Kelly FJ. Dietary antioxidants and environmental stress. *Procc Nutr Soc* 2004; 63(4): 579-585.
  21. Devereux G1, Turner SW, Craig LC, McNeill G, Martindale S, Harbour PJ, et al. Low maternal vitamin E intake during pregnancy is associated with asthma in 5-year-old children. *Am J Respir Crit Care Med* 2006; 174(5):499-507.
  22. Boots AW, Haenen GRMM, Bast A. Oxidant metabolism in chronic obstructive pulmonary disease. *Eur Respir J* 2003; 46: 14S-27S.
  23. Fogarty A, Lewis S, Weiss ST, Britton J. Dietary vitamin E, IgE concentrations, and atopy. *Lancet* 2000; 356(9241): 1573-4.
  24. Family food in 2005-06. UK Department of Environmental Farming and Rural Affairs Web site. <http://www.defra.gov.uk/evidence/statistics/foodfarm/food/familyfood/documents/familyfood-200506/pdf>. Posted 2007. Accessed December 9, 2010.
  25. National Food Survey datasets. UK Department of Environmental Farming and Rural Affairs Website. <http://www.defra.gov.uk/evidence/statistics/foodfarm/food/familyfood/nationalfoodsurvey/index.htm>. Updated March 2010. Accessed December 9, 2010.
  26. Henderson L, Irving K, Gregory J, Bates CJ, Prentice A, Perks J, Swan G, Farron M. The National Diet and Nutrition Survey: Adults aged 19 to 64 years. <http://www.food.gov.uk/multimedia/pdfs/ndnsv3.pdf>. Posted 2003. Accessed December 9, 2010.
  27. Ellwood P, Asher MI, Bjorksten B, Burr M, Pearce N, Robertson CF. Diet and asthma, allergic rhinoconjunctivitis and atopic eczema symptom prevalence: an ecological analysis of the International Study of Asthma and Allergies in Childhood (ISAAC) data. *ISAAC*

- Phase One Study Group. *Eur Respir J* 2001; 17(3):436–43.
28. Fogarty AW, Antoniak M, Venn AJ, Davies L, Goodwin A, Salfield N, Britton JR, Lewis SA. A natural experiment on the impact of fruit supplementation on asthma symptoms in children *Eur Respir J* 2009; 33(3): 481–5.
  29. Hatch GE. Asthma, inhaled oxidants, and dietary antioxidants. *Am J Clin Nutr* 1995; 61(Suppl.3): 625S–630S.
  30. Elsayed NM. Antioxidant mobilization in response to oxidative stress: a dynamic environmental-nutritional interaction. *Nutrition* 2001; 17(10): 828–834.
  31. Kolleck I, Sinha P, Rustow B. Vitamin E as an antioxidant of the lung: mechanisms of vitamin E delivery to alveolar type II cells. *Am J Respir Crit Care Med* 2002; 166(12Pt2): S62–66.
  32. Elsayed N, Mustafa M, Mead J. Increased vitamin E content in the lung after ozone exposure: a possible mobilization in response to oxidative, Bromberg stress. *Arch Biochem Biophys* 1990; 282(2): 263–269.
  33. Devereux G, Barker RN, Seaton A. Antenatal determinants of neonatal responses to allergens. *Clin Exp Allergy* 2002; 32(1): 43–50.
  34. Sienna-Monge JJ1, Ramirez-Aguilar M, Moreno-Macias H, Reyes-Ruiz NI, Del Río-Navarro BE, Ruiz-Navarro MX, et al. Antioxidant supplementation and nasal inflammatory responses among young asthmatics exposed to high levels of ozone. *Clin Exp Immunol* 2004; 138(2): 317–322.
  35. Brook RD, Brook JR, Rajagopalan S. Air pollution: the “Heart” of the problem. *Curr Hypertens Rep* 2003; 5(1): 32–39.
  36. Saxon A, Diaz-Sanchez D. Air pollution and allergy: you are what you breathe. *Nat Immunol* 2005; 6(3): 223–226.
  37. Grievink L, Smit HA, Ocké MC, van't Veer P, Kromhout D. Dietary intake of antioxidant (pro)-vitamins, respiratory symptoms and pulmonary function: the MORGEN study. *Thorax* 1998; 53(3): 166–71.
  38. Shaheen SO1, Sterne JA, Thompson RL, Songhurst CE, Margetts BM, Burney PG. Dietary antioxidants and asthma in adults: population-based case-control study. *Am. J. Respir. Crit. Care Med.* 2001; 164(10): 1823–8.
  39. Troisi RJ, Willett WC, Weiss ST, Trichopoulos D, Rosner B, Speizer FE. A prospective study of diet and adult-onset asthma. *Am. J. Respir. Crit. Care Med.* 1995; 151(1): 1401–8.
  40. Bodner C, Godden D, Brown K, Little J, Ross S, Seaton A. Antioxidant intake and adult-onset wheeze: a case-control study. Aberdeen WHEASE Study Group. *Eur. Respir. J.* 1999; 13(1): 22–30.
  41. Sackesen C, Ercan H, Dizdar E, Sover O, Gums P, Tosum BN, et al. A comprehensive evaluation of the enzymatic and nonenzymatic antioxidant systems in childhood asthma. *J Allergy Clinical Immunol* 2008; 122(1): 78–85.
  42. Kalayci O, Besler T, Kilinc K, Sekerel BE, Saraçlar Y. Serum levels of antioxidant vitamins (alpha tocopherol, beta carotene, and ascorbic acid) in children with bronchial asthma. *Turk J Pediatr* 2000; 42(1): 17-21.
  43. Misso NL, Brooks-Wildhaber J, Ray S, Vally H, Thompson PJ. Plasma concentration of dietary and non-dietary antioxidants are low in severe asthma. *Eur Respir J* 2005; 26(2): 257–64.
  44. Schock BC, Young IS, Brown V, Fitch PS, Shields MD, Ennis M. Antioxidants and oxidative stress in BAL fluid of a topic asthmatic children. *Pediatr Res* 2003; 53(3): 375-81.
  45. Nadeem A, Masood A, Siddiqui N. Oxidant-antioxidant imbalance in asthma: scientific evidence, epidemiological data and possible therapeutic options. *Ther Adv Respir Dis* 2008; 2(4): 215-35.
  46. Wood LG, Garg ML, Blake RJ, Simpson JL, Gibson PG. Oxidized vitamin E and glutathione as markers of clinical status in asthma. *Clin Nutr* 2008; 27(4): 579-86.
  47. Harik-Khan RI, Muller DC, Wise RA. Serum vitamin levels and the risk of asthma in children. *Am J Epidemiol* 2004; 159(4):351-7.

48. Gilliland FD, Berhane KT, Li YF, Gauderman WJ, McConnell R, Peters J. Children's lung function and antioxidant vitamin, fruit, juice, and vegetable intake. *Am J Epidemiol* 2003; 158(6): 576-84.
49. Hijazi N, Abalkhail B, Seaton A. Diet and childhood asthma in a society in transition: A study in urban and rural Saudi Arabia. *Thorax*. 2000; 55(9): 775-779.
50. Allen S, Britton J, Leonardi-Bee J. Association between antioxidant vitamins and asthma outcomes: Systematic review and meta-analysis. *Thorax*. 2009; 64: 610-619.
51. Chatzi L, Torrent M, Romieu I, Garcia-Esteban R, Ferrer C, Vioque J et al. Diet, wheeze, and atopy in school children in Menorca, Spain. *Pediatr Allergy Immunol*. 2007; 18(6): 480-485.
52. Chatzi L, Apostolaki G, Bibakis I, Skypala I, Bibaki-Liakou V, Tzanakis N, et al. Protective effect of fruits, vegetables and the Mediterranean diet on asthma and allergies among children in Crete. *Thorax*. 2007; 62(8): 677-683.
53. Forastiere F, Pistelli R, Sestini P, Fortes C, Renzoni E, Rusconi F, et al. Consumption of fresh fruit rich in vitamin C and wheezing symptoms in children. SIDRIA Collaborative Group, Italy (Italian Studies on Respiratory Disorders in Children and the Environment). *Thorax* 2000; 55(4): 283-288.
54. De Batlle J, Garcia-Aymerich J, Barraza-Villarreal A, Antó JM, Romieu I. Mediterranean diet is associated with reduced asthma and rhinitis in Mexican children. *Allergy* 2008; 63(10): 1310-1316.
55. Litonjua AA, Rifas-Shiman SL, Ly NP, Tantisira KG, Rich-Edwards JW, Camargo CA Jr, et al. Maternal antioxidant intake in pregnancy and wheezing illnesses in children at 2 y of age. *Am J Clin Nutr*. 2006; 84(4): 903-11.
56. Kelly FJ, Mudway I, Krishna MT, Holgate ST. The free radical basis of air pollution: focus on ozone. *Respir Med* 89(10): 647-656.
57. Mudway IS, Housley D, Eccles R, Richards RJ, Datta AK, Tetley TD, et al. Differential depletion of human respiratory tract antioxidants in response to ozone challenge. *Free Radic Res* 1996; 25: 499-513.
58. Schock B, Young IS, Fitch P, Brown V, Taylor R, Shields MD, et al. The effect of glass fronted fires on antioxidants in serum and BAL fluid from children. *Pediatr Res* 1998; 44(3): 455-455.
59. Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, et al. Effect of Mediterranean- Style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome. *JAMA* 2004; 292: 1440-6.
60. Cook DG, Carey IM, Whincup PH, Papacosta O, Chirico S, Bruckdorfer KR, et al. Effect of fresh fruit consumption on lung function and wheeze in children. *Thorax* 1997; 52(7): 628 - 33.
61. Farchi S, Forastiere F, Agabiti N, Corbo G, Pistelli R, Fortes C, et al. Dietary factors associated with wheezing and allergic rhinitis in children. *Eur Respir J* 2003; 22(5): 772- 80.
62. Forastiere F1, Pistelli R, Sestini P, Fortes C, Renzoni E, Rusconi F, et al. Consumption of fresh fruit rich in vitamin C and wheezing symptoms in children. SIDRIA Collaborative Group, Italy (Italian Studies on Respiratory Disorders in Children and the Environment). *Thorax* 2000;55(4): 283-8.
63. Nja F, Nystad W, Lodrup Carlsen KC, Hetlevik O, Carlsen KH. Effects of early intake of fruit or vegetables in relation to later asthma and allergic sensitization in school-age children. *Acta Paediatr* 2005; 94(2): 147-54.
64. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Consumption of vegetables, fruit, and antioxidants during pregnancy and wheeze and eczema in infants. *Allergy*. 2010; 65(6):758-765.
65. Greenough A, Shaheen SO, Shennan A, Seed PT, Poston L. Respiratory outcomes in early childhood following antenatal vitamin C and E supplementation. *Thorax*. 2010; 65(11): 998-1003.
61. Devereaux G, Barker RN, Seaton A. Antenatal determinants of neonatal responses



- to allergens. *Clin Exp Allergy* 2002; 32(1): 43–50.
66. Martindale S, McNeill G, Devereux G, Campbell D, Russell G, Seaton A. Antioxidant intake in pregnancy in relation to wheeze and eczema in the first two years of life. *Am J Respir Crit Care Med* 2005; 171(2): 121–8.
67. Litonjua, A. A., Tantisira, K. G., Finn, P. W., Schaub, B., Schroeter, C., Perkins, D. L. Maternal antioxidant intake during pregnancy and cord blood lymphoproliferative responses. *Am J Respir Crit Care Med* 2004; 169: A501.
68. Clark J, Craig L, McNeill G, Smith N, Norrie J, Devereux GA. A novel dietary intervention to optimize vitamin E intake of pregnant women to 15 mg/day. *J Acad Nutr Diet.* 2012 Feb;112(2):297-301.
69. Devereux G, Turner SW, Craig LCA, McNeill G, Martindale S, Harbour PJ, et al. Reduced maternal vitamin E intake during pregnancy is associated with asthma in 5-year-old children. *Am J Respir Crit Care Med.* 2006;174: 499–507.
70. Litonjua AA, Rifas-Shiman S, Ly NP, Tantisira KG, Rich-Edwards JW, Weiss ST, et al. Maternal antioxidant intake in pregnancy and wheezing illnesses in children at 2 y of age. *Am J Clinical Nutr.* 2006;84:903–911.
71. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Consumption of vegetables, fruit, and antioxidants during pregnancy and wheeze and eczema in infants. *Allergy.* 2010;65:758–765.