

Accepted Manuscript

Accepted Manuscript (Uncorrected Proof)

Title: Targeting Copper Dyshomeostasis as a Pathophysiological Basis of Childhood Obesity - the Latest Facts

Running Title: The Relationship between Copper Status and Childhood Obesity

Authors: Marina Jakšić^{1,*}, Milica Martinović², Mirjana Nedović Vuković³

1. *Department of Laboratory Diagnostics, Institute for Children's Diseases, Clinical Center of Montenegro, Podgorica, Montenegro.*
2. *Department of Pathological Physiology, Medical Faculty, University of Montenegro, Podgorica, Montenegro.*
3. *Center for Health System Evidence and Research in Public Health, Department of Health Statistics, Institute for Public Health of Montenegro, Podgorica, Montenegro.*

***Corresponding Author:** Marina Jakšić, Department of Laboratory Diagnostics, Institute for Children's Diseases, Clinical Center of Montenegro, Podgorica, Montenegro. E-mail: marinaj@ucg.ac.me

To appear in: **Journal of Pediatrics Review**

Received date: 2023/12/05

Revised date: 2024/01/21

Accepted date: 2024/02/05

This is a “Just Accepted” manuscript, which has been examined by the peer-review process and has been accepted for publication. A “Just Accepted” manuscript is published online shortly after its acceptance, which is prior to technical editing and formatting and author proofing. Journal of Pediatrics Review provides “Just Accepted” as an optional and free service which allows authors to make their results available to the research community as soon as possible after acceptance. After a manuscript has been technically edited and formatted, it will be removed from the “Just Accepted” web site and published as a published article. Please note that technical editing may introduce minor changes to the manuscript text and/or graphics which may affect the content, and all legal disclaimers that apply to the journal pertain.

Please cite this article as:

Jakšić M, Martinović M, Nedović Vuković M. Targeting Copper Dyshomeostasis as a Pathophysiological Basis of Childhood Obesity - the Latest Facts. Journal of Pediatrics Review. Forthcoming 2024.

Abstract

Context: Childhood pre-obesity and obesity rates have been rapidly growing worldwide over the past decades. Copper homeostasis is gaining increasing attention in the physiopathology of obesity. Strong evidences indicate that a disturbance of copper homeostasis plays an important role in the development of obesity and its related comorbidities. Under physiological conditions, copper plays a significant role in regulatory, immunologic, and antioxidant functions resulting in protection against inflammation and oxidative stress, and consequently against the known comorbidities of obesity. Nevertheless, despite the growing body of research, information about copper status in obesity, particularly in childhood obesity, is scarce.

Evidence Acquisition: The aim of this brief narrative review is to examine the latest data published in the last five years using databases such as PubMed, Scopus, Unpaywall (COBISS), and EBSCO to emphasize the major current findings in research related to this topic.

Results and Conclusions: The most recent studies have yielded strong evidence in support of altered copper status in childhood obesity, nevertheless, further studies are needed to clarify the role of copper in the physiopathology of childhood obesity.

Keywords: Childhood obesity, Childhood overweight, Serum copper, Copper homeostasis

1. Context

According to the World Health Organization (WHO), overweight and obesity are defined as abnormal or excessive fat accumulation that may impair health and have become one of the most significant epidemics of the 21st century (1).

Alerting data show that overweight and obesity are increasing rapidly worldwide; overall, about 13% of the world's adult population and 340 million children and adolescents aged 5-19 were overweight or obese in 2016 (1). According to current trends, 1 in 5 adults worldwide is expected to be affected by obesity by 2025. These data clearly show that obesity is an alarmingly increasing global public health issue (2).

An anthropometric indicator that is commonly used as a screening tool for overweight and obesity in adults is body mass index (BMI, kg/m²). BMI categories for children and teenagers are based on sex- and age-specific BMI percentiles (Table 1) (3):

Table 1. Child BMI categories and their corresponding sex- and age-specific BMI percentiles

BMI Category	BMI Range
Underweight	Less than the 5 th percentile
Healthy Weight	5 th percentile to less than the 85 th percentile
Overweight	85 th percentile to less than the 95 th percentile
Obesity	95 th percentile or greater

Overweight and obesity are influenced by genetics, biology, psychosocial factors, and health behaviors. Furthermore, the primary risk factors for obesity - poor diet and physical inactivity - are among the top causes of preventable youth mortality, chronic disease, and economic health burden (4).

It is a well-known fact that obesity increases the risk of non-communicable chronic diseases and cardiometabolic risks (5) in children, such as metabolic syndrome, hypertension (6,7), dyslipidemia (8), insulin resistance and diabetes mellitus type 2 (9) nonalcoholic fatty liver disease (NAFLD), (10), certain cancer (11), autoimmune diseases, as well as depression and cognitive disturbances that may contribute to poor school results (12). Children who are overweight or obese are more prone to become obese adults and develop comorbidities at an earlier age than children who are regarded to be of healthy weight. There is also an increased risk of premature death and impairment in later life (13).

Although obesity is largely preventable, it is difficult to treat given its multifactorial nature. Weight management strategies include dietary modification, increased physical activity, and medical or surgical management for severe and morbid obesity (14). The roles of oligo-elements in obesity have recently attracted increased attention due to their oxidant or antioxidant actions, as well as their impacts on insulin/glucose and lipid metabolism, which may be linked to obesity (15). Studies have shown that metabolic diseases including obesity are closely related to systemic inflammation, oxidative stress, and disorders of copper and iron metabolism (16). Additionally, it is shown that serum copper might increase as an acute-phase response in a variety of inflammatory conditions, including obesity-induced low-grade chronic inflammation (17). The investigation of sophisticated molecular mechanisms that could potentially become suitable therapeutic targets remains one of the most important research aspirations worldwide, and this is exactly where the suggested link between copper metabolism and obesity pathophysiology is in line.

2. Evidence Acquisition

English-language relevant literature from the past 5 years was considered, including studies from a broad range of methodologies: original papers, meta-analyses, clinical trials, and reviews. Letters and case reports were not included. Research terms adopted, alone and/or combined, were obesity, obese, overweight, adiposity, adolescents, children, trace elements, essential elements, copper, hypertension, and dyslipidemia. The databases PubMed, Scopus, Unpaywall (COBISS), and EBSCO were used for the literature research.

3. Altered copper metabolism and childhood obesity – is there a more comprehensible link?

Copper is a redox-active substance that participates in many biological processes and may be involved in a multitude of pathogenic events, including insulin resistance, inflammation, oxidative stress, abnormal glucose metabolism, and dyslipidemia (18) (Figure 1) #

Under physiological conditions, as cofactors for enzymes, copper ions are required for processes such as cellular energy production, modulation of synaptic transmission, iron metabolism, angiogenesis, inflammatory response, and antioxidant defense (19,20). Paradoxically, an excessive level of serum copper can cause an increase in reactive oxygen species (ROS) in the organism, which is in agreement with the increased oxidative stress found in childhood obesity (19). The copper status of obese individuals is usually evaluated by measuring levels of copper and ceruloplasmin in serum, adipose, and hepatic tissues (21).

Even though reported in some studies, information about copper homeostasis in human obesity is limited (20). Accordingly, information about copper homeostasis in childhood obesity is even more scarce. Nevertheless, the serum copper alterations have been proposed to have associations

among children with obesity (18, 22, 23). Chandrashekhar et al. (2023) established a correlation between serum copper and obesity in 51 adolescents aged 13-16 years with BMI < 95th percentile, demonstrating an excess of serum copper in obese individuals (24). In a study conducted by Castillo-Valenzuela et al. (2023) which included 1235 normal-weight and obese participants, it was found that children between the ages of 4 and 14 had deficiencies in vitamin D, as well as copper, calcium, and zinc to a lesser extent (25). Many studies have documented elevated circulating copper levels in obese children (18, 23, 25, 26, 27, 28, 29, 30). In addition, Meggyesy et al. (2020) demonstrated in a diet-induced obesity mouse model that copper ionophores administered daily per os, augmented hepatic copper and decreased mouse body weight, which established copper ionophores as a potential class of antiobesity agents (31). Wang et al (2023) hypothesized that the altered lipid profiles associated with copper overload may contribute to obesity-related systemic inflammation and oxidative stress (15). However, some studies demonstrated that the serum copper levels were significantly lower in children with obesity as compared to controls (22). Then again, some authors reported no significant difference in serum copper levels between obese and normal-weight children (32) or any relation between metabolic syndrome and plasma copper in investigated children (33) Gonzalez Domingez et al. (2023) studied an observational cohort comprising 46 prepubertal and 48 pubertal children with obesity, and have demonstrated that non-obese children going through puberty had better control over inflammation and oxidative stress. Additionally, they had higher levels of essential elements that are involved in the antioxidant system and metabolic control (such as zinc, molybdenum, selenium, and manganese). Inversely, total copper and free iron were found to be reduced in their blood (34). These results clearly show that copper is a significant modulator of the adipocyte metabolism; nevertheless, the fundamental mechanism of copper's contribution to fat cell pathophysiology is still mostly unclear (35).

Together with some other trace elements, data show that copper plays an indispensable role in cardiovascular protection (36) and cholesterol modulation (37), two conditions that are significantly affected by pre-obesity and obesity even in childhood. Some studies show that copper deficiency has been suggested to contribute to cardiovascular disease, which is one of the best-known consequences of obesity. Accordingly, Tong et al. (2022) found that childhood hypertension was negatively correlated with the dietary intake of copper, and was found positively correlated with triglyceride level ≥ 1.69 mmol/L, low-density lipoprotein cholesterol level ≥ 2.84 mmol/L, BMI z-score and central obesity (38). As opposed to these findings, high serum copper contents have been reported to be independent risk factors and biomarkers of cardiovascular diseases as well (27, 36). Additionally, a National Health and Nutrition Examination Survey (2011-2016) in the United States of America suggested that high serum copper concentrations were significantly associated with elevated blood pressure in US children and adolescents (39).

The results of the cross-sectional study which included 3982 children and adolescents who participated in the US National Health and Nutrition Examination Survey 1999-2006 showed that higher dietary copper intake increases the prevalence of hypertriglyceridemia, especially among US adolescents with a BMI ≥ 23 kg/m² (40). A partial underlying mechanism was suggested by Zhong et al. (2022), who found that copper excess activated oxidative stress and autophagy, up-regulated lipogenesis and lipid metabolism (41). Similar findings were reported by González-Domínguez et al. (2022), who showed that abnormalities related to trace elements including high serum copper were generally more prevalent in obese children and associated with dyslipidemia, inflammation, oxidative stress, and improper glucose metabolism (18). Nevertheless, Blades et al. (2021) proposed a different copper and lipid metabolism interrelationship, claiming that the

increased cellular copper downregulates lipids and lipogenic genes, and vice versa. In this study, obesity and increased dietary cholesterol corresponded with decreased tissue copper (37), which is an interesting and distinct outcome from the majority of the other research given in this review.

4. Conclusion

After conducting a thorough examination of the latest scientific literature on copper metabolism and childhood obesity, our findings suggest that there is a significant association between childhood obesity and elevated serum copper concentrations, as evidenced by the majority (but not all) of studies conducted in this area. However, while this correlation is evident, further research is needed to determine the exact causal relationship between the two, as well as to uncover the underlying cellular and subcellular mechanisms that drive this relationship. The existing literature presents scarce explanations for the observed association. Some researchers have proposed that copper may contribute to the development of obesity by promoting lipid synthesis and storage, while others have suggested that copper may exacerbate inflammation in adipose tissue, leading to insulin resistance and other metabolic dysfunctions associated with obesity. Despite these potential explanations, there remains a significant gap in our understanding of the role of copper in childhood obesity. As such, future research must focus on elucidating the causal relationship between copper and obesity, as well as the underlying biological mechanisms involved.

Conflict of interest statement: The authors declare no conflict of interest.

Authorship Contribution: Conceptualization: M.J; Methodology: M.J; Writing – Original Draft Preparation: M.J., M.M., M.N.V; Writing – Review & Editing, M.J., M.M;

References:

1. WHO Obesity and overweight. Available at: <https://www.who.int/news-room/factsheets/detail/obesity-and-overweight> (cited September 2023)
2. World Obesity. The obesity pandemic. Available at: <https://www.worldobesity.org/news/the-time-to-act-is-now-accelerating-action-on-obesity-during-covid-19> (cited September 2023)
3. Centers for Disease Control and Prevention. Defining Child BMI Categories. Available at: <https://www.cdc.gov/obesity/basics/childhooddefining.html> (cited Jan, 2024)
4. Smith JD, Fu E, Kobayashi MA. Prevention and Management of Childhood Obesity and Its Psychological and Health Comorbidities. *Annu Rev Clin Psychol*. 2020 May 7;16:351-378. doi: 10.1146/annurev-clinpsy-100219-060201
5. Martinović M, Belojević G, Jakšić M, Kavarić N, Klisić A. Cardiometabolic Risk Among Montenegrin Urban Children In Relation To Obesity And Gender. *Acta Clin Croat*. 2021;60(1):3-9. doi:10.20471/acc.2021.60.01.01
6. Litwin M, Kułaga Z. Obesity, metabolic syndrome, and primary hypertension. *Pediatr Nephrol*. 2021;36(4):825-837. doi: 10.1007/s00467-020-04579-3
7. Martinovic M, Belojevic G, Evans GW, Kavaric N, Asanin B, Pantovic S, Jaksic M, Boljevic J. Hypertension and correlates among Montenegrin schoolchildren-a cross-sectional study. *Public Health*. 2017 Jun;147:15-19. doi: 10.1016/j.puhe.2017.02.007. Epub 2017 Mar 6. PMID: 28404491.
8. Vekic J, Zeljkovic A, Stefanovic A, Jelic-Ivanovic Z, Spasojevic-Kalimanovska V. Obesity and dyslipidemia. *Metabolism*. 2019;92:71-81. doi: 10.1016/j.metabol.2018.11.005
9. Rendell MS. Obesity and diabetes: the final frontier. *Expert Rev Endocrinol Metab*. 2023;18(1):81-94. doi:10.1080/17446651.2023.2168643

10. Younossi Z, Tacke F, Arrese M, Chander Sharma B, Mostafa I, Bugianesi E, Wai-Sun Wong V, Yilmaz Y, George J, Fan J, Vos MB. Global Perspectives on Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. *Hepatology*. 2019;69(6):2672-2682. doi:10.1002/hep.30251
11. Kim DS, Scherer PE. Obesity, Diabetes, and Increased Cancer Progression. *Diabetes Metab J*. 2021;45(6):799-812. doi:10.4093/dmj.2021.007
12. Lindberg L, Persson M, Danielsson P, Hagman E, Marcus C. Obesity in childhood, socioeconomic status, and completion of 12 or more school years: a prospective cohort study. *BMJ Open*. 2021;11:e040432
13. The Lancet Diabetes Endocrinology. Childhood obesity: a growing pandemic. *Lancet Diabetes Endocrinol*. 2022;10(1):1. doi: 10.1016/S2213-8587(21)00314-4
14. Jebeile H, Kelly AS, O'Malley G, Baur LA. Obesity in children and adolescents: epidemiology, causes, assessment, and management. *Lancet Diabetes Endocrinol*. 2022;10(5):351-365. doi:10.1016/S2213-8587(22)00047-X
15. Wang L, Liu W, Bi S, Zhou L, Li L. Association between minerals intake and childhood obesity: A cross-sectional study of the NHANES database in 2007-2014. *PLoS One*. 2023;18(12):e0295765. doi: 10.1371/journal.pone.0295765
16. Liu Z, Wang M, Zhang C, Zhou S, Ji G. Molecular Functions of Ceruloplasmin in Metabolic Disease Pathology. *Diabetes Metab Syndr Obes*. 2022 Mar 3;15:695-711. doi: 10.2147/DMSO.S346648
17. Escobedo-Monge MF, Barrado E, Parodi-Román J, Escobedo-Monge MA, Torres-Hinojal MC, Marugán-Miguelsanz JM. Copper and Copper/Zn Ratio in a Series of Children with Chronic Diseases: A Cross-Sectional Study. *Nutrients*. 2021;13(10):3578. doi: 10.3390/nu13103578
18. González-Domínguez Á, Millán-Martínez M, Domínguez-Riscart J, Mateos RM, Lechuga-Sancho AM, González-Domínguez R. Altered Metal Homeostasis Associates with Inflammation,

- Oxidative Stress, Impaired Glucose Metabolism, and Dyslipidemia in the Crosstalk between Childhood Obesity and Insulin Resistance. *Antioxidants (Basel)*. 2022;11(12):2439. Published 2022 Dec 10. doi:10.3390/antiox11122439
19. Jomova K, Makova M, Alomar SY, Alwasel SH, Nepovimova E, Kuca K, Rhodes CJ, Valko M. Essential metals in health and disease. *Chem Biol Interact*. 2022;367:110173. doi:10.1016/j.cbi.2022.110173
20. Yang H, Liu CN, Wolf RM, Ralle M, Dev S, Pierson H, Askin F, Steele KE, Magnuson TH, Schweitzer MA, Wong GW, Lutsenko S. Obesity is associated with copper elevation in serum and tissues. *Metallomics*. 2019;11(8):1363-1371. doi:10.1039/c9mt00148d
21. Calcaterra V, Verduci E, Milanta C, Agostinelli M, Todisco CF, Bona F, Dolor J, La Mendola A, Tosi M, Zuccotti G. Micronutrient Deficiency in Children and Adolescents with Obesity—A Narrative Review. *Children*. 2023; 10(4):695. <https://doi.org/10.3390/children10040695>
22. Vivek SM, Dayal D, Khaiwal R, Bharti B, Bhalla A, Singh S, Kaur H, Attri SV. Low serum copper and zinc concentrations in North Indian children with overweight and obesity. *Pediatr Endocrinol Diabetes Metab*. 2020;26(2):79-83. English. doi: 10.5114/pedim.2020.95627. PMID: 32564583
23. Ge W, Liu W, Liu G. The relationships between serum copper levels and overweight/total obesity and central obesity in children and adolescents aged 6-18 years (published online ahead of print, 2020 May 20). *J Trace Elem Med Biol*. 2020;61:126557. doi:10.1016/j.jtemb.2020.126557
24. Chandrashekhar A, Bembalgi S, Malebennur SK. Serum Zinc and Copper Levels in Obese Adolescents, *TheScientificTemper*, 2023;14(3):10551059. Doi10.58414/SCIENTIFICTEMPER.2023.14.3.75

25. Castillo-Valenzuela O, Duarte L, Arredondo M, Iñiguez G, Villarroel L, Pérez-Bravo F Childhood Obesity and Plasma Micronutrient Deficit of Chilean Children between 4 and 14 Years Old, *Nutrients* 2023; 15(7), 1707. doi.org/10.3390/nu15071707
26. Thillan K, Lanerolle P, Thoradeniya T, Samaranayake D, Chandrajith R, Wickramasinghe P. Micronutrient status and associated factors of adiposity in primary school children with normal and high body fat in Colombo municipal area, Sri Lanka. *BMC Pediatr.*2021;21(1):14. doi:10.1186/s12887-020-02473-3
27. Leone N, Courbon D, Ducimetiere P, Zureik M. Zinc, copper, and magnesium and risks for all-cause, cancer, and cardiovascular mortality. *Epidemiology.* 2006;17(3):308-14. doi: 10.1097/01.ede.0000209454.41466.b7
28. Gu K, Li X, Xiang W, Jiang X. The Relationship Between Serum Copper and Overweight/Obesity: a Meta-analysis. *Biol Trace Elem Res.* 2020;194(2):336-347. doi:10.1007/s12011-019-01803-6
29. Vazquez-Moreno M, Sandoval-Castillo M, Rios-Lugo MJ, Klünder-Klünder M, Cruz M, Martínez-Navarro I, Romero-Guzmán ET, Victoria-Campos CI, Vilchis-Gil J, Hernández-Mendoza H. Overweight and Obesity Are Positively Associated with Serum Copper Levels in Mexican Schoolchildren. *Biol Trace Elem Res.* 2023;201(6):2744-2749. doi:10.1007/s12011-022-03383-4
30. Du M, Qiu M, Qian Y, Wang T, Chen X. Serum Copper/Zinc Ratio in Overweight and Obese Children: a Cross-Sectional Study. *Biol Trace Elem Res.* 2023; doi: 10.1007/s12011-023-03790-1 Epub ahead of print
31. Meggyesy PM, Masaldan S, Clatworthy SAS, Volitakis I, Eyckens DJ, Aston-Mourney K, Cater MA. Copper Ionophores as Novel Antiobesity Therapeutics. *Molecules.* 2020;25(21):4957. doi: 10.3390/molecules25214957

32. Jaksic M, Martinovic M, Gligorovic-Barhanovic N, Vujacic A, Djurovic D, Nedovic-Vukovic M. Association between inflammation, oxidative stress, vitamin D, copper and zinc with pre-obesity and obesity in school children from the city of Podgorica, Montenegro. *J Pediatr Endocrinol Metab.* 2019;32(9):951-957. doi:10.1515/jpem-2019-0086
33. Villatoro-Santos CR, Ramirez-Zea M, Villamor E; Nine Mesoamerican Countries Metabolic Syndrome (NiMeCoMeS) Study Group. Plasma Copper and Metabolic Syndrome in Mesoamerican Children and Their Parents. *Biol Trace Elem Res.* 2024; doi: 10.1007/s12011-024-04069-9. Epub ahead of print
34. González-Domínguez A, Domínguez Riscart J, Millán-Martínez M, María Mateos-Bernal RM, Lechuga-Sancho AM, González-Domínguez R. Trace elements as potential modulators of puberty-induced amelioration of oxidative stress and inflammation in childhood obesity, *BioFactors* 2023; 49(4): 820-830. doi.10.1002/biof.1946
35. Yang H, Ralle M, Wolfgang MJ, Dhawan N, Burkhead JL, Rodriguez S, Kaplan JH, Wong GW, Haughey N, Lutsenko S. Copper-dependent amino oxidase 3 governs selection of metabolic fuels in adipocytes. *PLoS Biol.* 2018;16(9):e2006519. doi: 10.1371/journal.pbio.2006519
36. Shi Y, Zou Y, Shen Z, Xiong Y, Zhang W, Liu C, Chen S. Trace Elements, PPARs, and Metabolic Syndrome. *Int J Mol Sci.* 2020;21(7):2612. doi: 10.3390/ijms21072612
37. Blades B, Ayton S, Hung YH, Bush AI, La Fontaine S. Copper and lipid metabolism: A reciprocal relationship. *Biochim Biophys Acta Gen Subj.* 2021;1865(11):129979. doi:10.1016/j.bbagen.2021.129979
38. Tong J, An X, Zhao L, Qu P, Tang X, Chen M, Liang X. Combining multiaspect factors to predict the risk of childhood hypertension incidence. *J Clin Hypertens.* 2022; 24: 1015–1025. <https://doi.org/10.1111/jch.14544>

39. Liu C, Liao Y, Zhu Z, Yang L, Zhang Q, Li L. The association between serum copper concentrations and elevated blood pressure in US children and adolescents: National Health and Nutrition Examination Survey 2011-2016. *BMC Cardiovasc Disord.* 2021;21(1):57. doi:10.1186/s12872-021-01880-3
40. Shi Y, Hu H, Wu Z, Wu J, Chen Z, Cheng X, Li P. Associations between dietary copper intake and hypertriglyceridemia among children and adolescents in the US. *Nutr Metab Cardiovasc Dis.* 2023;33(4):809-816. doi:10.1016/j.numecd.2023.01.020
41. Zhong CC, Zhao T, Hogstrand C, Chen F, Song CC, Luo Z. Copper (Cu) induced changes of lipid metabolism through oxidative stress-mediated autophagy and Nrf2/PPAR γ pathways. *J Nutr Biochem.* 2022;100:108883. doi:10.1016/j.jnutbio.2021.108883